

Fig.6.





Fig.7.



10

 $V_{GS}(V)$

15

5

0

-10

-5

20

25

Fig.8.

- --- On And Off Current With No Field Enhancement
- → Off Current With Field Enhancement (Model By Hurkx et al)
- -x- Off Current With Field Enhancement (Present Invention)




Fig.9.

$$\Gamma_n^{Coul} = \frac{\Delta E_n}{kT} \int_{\Delta E_{fp}}^{\infty} \exp \left\{ \frac{\Delta E_n}{kT} u - K_n u^{\frac{3}{2}} \left[1 - \left(\frac{\Delta E_{fp}}{u \Delta E_n} \right)^{\frac{5}{3}} \right] \right\} du$$
 (1)

$$\Gamma_n^{Coul} = \frac{\Delta E_n}{kT} \int_{\Delta E_{fp}}^{1} \exp\left\{ \frac{\Delta E_n}{kT} u - K_n u^{\frac{3}{2}} + K_n \left(\frac{\Delta E_{fp}}{u\Delta E_n} \right)^{\frac{5}{3}} u^{-\frac{1}{6}} \right\} du.$$
 (2)

$$A = \frac{\Delta E_n}{kT}, B = K_n, C = \frac{\Delta E_{fp}}{\Delta E_n}, D = BC^{\frac{5}{3}}.$$
 (3)

$$\Gamma_n^{Coul} = A \int_c^l \exp\left\{Au - Bu^{\frac{3}{2}} + Du^{-\frac{1}{6}}\right\} du. \tag{4}$$

$$\Gamma_n^{Coul} = A \int_c^l \exp[f(u)] du.$$
 (5)

$$f(u) = Au - Bu^{\frac{3}{2}} + Du^{-\frac{1}{6}}.$$
 (6)

$$f(u) \approx f(v) + f'(v)(u - v) + \frac{f''(v)}{2}(u - v)^2,$$
 (7)

Fig.9(cont.a)

$$f(v) = Av - Bv^{\frac{3}{2}} + Dv^{-\frac{1}{6}}.$$
 (8)

$$f'(v) = A - \frac{3}{2}Bv^{\frac{1}{2}} - \frac{1}{6}Dv^{-\frac{7}{6}},\tag{9}$$

$$f''(v) = -\frac{3}{4}Bv^{-\frac{1}{2}} + \frac{7}{36}Dv^{-\frac{13}{6}}.$$
 (10)

$$f(u) \approx \frac{f''(v)}{2}u^2 + [f'(v) - vf''(v)]u + \left[v^2 \frac{f''(v)}{2} - vf'(v) + f(v)\right]. \tag{11}$$

$$f(u) \approx f''(v) u + [f'(v) - vf''(v)] = 0,$$

$$u_{\text{max}} = \frac{f'(v) - vf''(v)}{f''(v)}.$$

$$f(u) \approx -(AIu^2 + AIIu + AIII)$$
 (12)

$$AI = -\frac{f''(v)}{2}. ag{13}$$

$$A\Pi = -[f'(v) - vf''(v)] \tag{14}$$

$$AIII = -\left[v^2 \frac{f''(v)}{2} - vf'(v) + f(v)\right]. \tag{15}$$

$$f(u) \approx -AI \left[\left(u + \frac{AII}{2AI} \right)^2 + \left(\frac{AIII}{AI} - \left(\frac{AII}{2AI} \right)^2 \right) \right]. \tag{16}$$

$$\Gamma_n^{Coul} = A \int_{c}^{l} \exp[f(u)] du$$
 (17)

Fig.9(cont.b)

$$\Gamma_n^{Coul} = A \exp \left[AI \left(\frac{AII}{AI} - \left(\frac{AII}{2AI} \right)^2 \right) \right] \int_C^I \exp \left[\sqrt{AI} \left(u + \frac{AII}{2AI} \right) \right]^2 du.$$
 (18)

$$t = \sqrt{AI} \left(u + \frac{AII}{2AI} \right), \tag{19}$$

$$u = C, \ t_l = \sqrt{AI} \left(C + \frac{AII}{2AI} \right), \tag{20}$$

$$u = I, t_u = \sqrt{AI} \left(1 + \frac{AII}{2AI} \right), \tag{21}$$

$$du = \frac{1}{\sqrt{AI}} dt. ag{22}$$

$$\Gamma_n^{Coul} = \frac{A}{\sqrt{AI}} \exp \left[AI \left(\frac{AII}{AI} - \left(\frac{AII}{2AI} \right)^2 \right) \right]_{t_I}^{t_U} e^{-t^2} dt.$$
 (23)

$$\int_{t_{I}}^{t_{U}} e^{-t^{2}} dt = \frac{\sqrt{\pi}}{2} \left[erf(t_{U}) - erf(t_{I}) \right].$$
 (24)

$$erf(x) = 1 - (a_1t + a_2t^2 + a_3t^3 + a_4t^4 + a_5t^5)e^{-x^2}$$
,

$$t=\frac{1}{1+px}\,,$$

$$a_1 = 0.254829592;$$

$$a_2 = -0.284496736$$
;

$$a_3 = 1.421413741;$$

$$a_4 = -1.453152027;$$

$$a_5 = 1.061405429;$$

p = 0.3275911; (25)

Fig.9(cont.c)

$$\Gamma_n^{Coul} = \frac{A}{2} \sqrt{\frac{\pi}{I}} \left(\frac{a_1}{(1+pt_0)} + \frac{a_2}{(1+pt_0)^2} + \frac{a_3}{(1+pt_0)^3} + \frac{a_4}{(1+pt_0)^4} + \frac{a_5}{(1+pt_0)^5} \right) \exp(-C^2 AI - CAII - AIII)$$

$$- \frac{A}{2} \sqrt{\frac{\pi}{I}} \left(\frac{a_1}{(1+pt_0)} + \frac{a_2}{(1+pt_0)^2} + \frac{a_3}{(1+pt_0)^3} + \frac{a_4}{(1+pt_0)^4} + \frac{a_5}{(1+pt_0)^5} \right) \exp(-AI - AIII)$$

$$AI = -\frac{f''(v)}{2}, AII = -[f'(v) - vf''(v)], AIII = -[v^2 \frac{f''(v)}{2} - vf'(v) + f(v)],$$

$$t_{l} = \sqrt{AI} \left(C + \frac{AII}{2AI} \right), \ t_{u} = \sqrt{AI} \left(1 + \frac{AII}{2AI} \right),$$

$$f(v) = Av - Bv^{\frac{3}{2}} + Dv^{-\frac{1}{6}},$$

$$f'(v) = A - \frac{3}{2} Bv^{\frac{1}{2}} - \frac{1}{6} Dv^{-\frac{7}{6}},$$

$$f''(v) = -\frac{3}{4} Bv^{-\frac{1}{2}} + \frac{7}{36} Dv^{-\frac{13}{6}},$$

Fig.9(cont.d)

$$A = \frac{\Delta E_n}{kT}$$
, $B = K_n$, $C = \frac{\Delta E_{fp}}{\Delta E_n}$, $D = BC^{\frac{5}{3}}$.

$$v = C(\text{for } u_{max} < C, \text{ case 1}),$$

$$v = u_{max}$$
 (for $C < u_{max} < I$, case 2),

$$v = I(\text{for } u_{max} \ge 1, \text{ case } 3),$$

$$u_{\text{max}} = \frac{f'(v) - vf''(v)}{f''(v)}$$
 for $v = \frac{C+1}{2}$

$$\Gamma_n^{Coul} = \frac{A}{2} \sqrt{\frac{\pi}{AI}} \left(\frac{a_1}{(1+pt_l)} + \frac{a_2}{(1+pt_l)^2} + \frac{a_3}{(1+pt_l)^3} + \frac{a_4}{(1+pt_l)^4} + \frac{a_5}{(1+pt_l)^5} \right) \exp(-C^2AI - CAII - AIII)$$

$$- \frac{A}{2} \sqrt{\frac{\pi}{AI}} \left(\frac{a_1}{(1+pt_u)} + \frac{a_2}{(1+pt_u)^2} + \frac{a_3}{(1+pt_u)^3} + \frac{a_4}{(1+pt_u)^4} + \frac{a_5}{(1+pt_u)^5} \right) \exp(-AII - AIII - AIII)$$

$$\pm \frac{A}{2} \sqrt{\frac{\pi}{AI}} \exp\left(-AIII + \frac{AIII}{4AII}\right)$$